

Covariance and Total Expectation Intro

Covariance: measure of the relationship between two RVs

$$\text{cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])] = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y].$$

The sign of $\text{cov}(X, Y)$ illustrates how X and Y are related; a positive value means that X and Y tend to increase and decrease together, while a negative value means that X increases as Y decreases (and vice versa). A covariance of zero means that the two random variables are uncorrelated—there is no relationship between them.

Properties: for random variables X, Y, Z and constant a ,

- $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2\text{cov}(X, Y)$
- $\text{cov}(X, X) = \text{Var}(X)$
- $\text{cov}(X, Y) = \text{cov}(Y, X)$
- Bilinearity: $\text{cov}(X + Y, Z) = \text{cov}(X, Z) + \text{cov}(Y, Z)$ and $\text{cov}(aX, Y) = a\text{cov}(X, Y)$

Conditional Expectation: When we want to find the expectation of a random variable X conditioned on an event A , we use the following formula:

$$\mathbb{E}[X | A] = \sum_x x \cdot \mathbb{P}[(X = x) | A].$$

This is an application of the definition of expectation. We still consider all values of X but reweigh them based on their probability of occurring together with A .

Total Expectation: For any random variable X and events A_1, A_2, \dots, A_n that partition the sample space Ω ,

$$\mathbb{E}[X] = \sum_{i=1}^n \mathbb{E}[X | A_i] \mathbb{P}[A_i].$$

We can think of this as splitting the sample space into partitions (events) and looking at the expectation of X in each partition, weighted by the probability of that event occurring.

1 Covariance

Note 16

- (a) We have a bag of 5 red and 5 blue balls. We take two balls uniformly at random from the bag without replacement. Let X_1 and X_2 be indicator random variables for the events of the first and second ball being red, respectively. What is $\text{cov}(X_1, X_2)$? Recall that $\text{cov}(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$.
- (b) Now, we have two bags A and B, with 5 red and 5 blue balls each. Draw a ball uniformly at random from A, record its color, and then place it in B. Then draw a ball uniformly at random from B and record its color. Let X_1 and X_2 be indicator random variables for the events of the first and second draws being red, respectively. What is $\text{cov}(X_1, X_2)$?

2 Number Game

Note 20

Sinho and Vrettos are playing a game where they each choose an integer uniformly at random from $[0, 100]$, then whoever has the larger number wins (in the event of a tie, they replay). However, Vrettos doesn't like losing, so he's rigged his random number generator such that it instead picks randomly from the integers between Sinho's number and 100. Let S be Sinho's number and V be Vrettos' number.

(a) What is $\mathbb{E}[S]$?

(b) What is $\mathbb{E}[V \mid S = s]$, where s is any constant such that $0 \leq s \leq 100$?

(c) What is $\mathbb{E}[V]$?

3 Dice Games

Note 20

Suppose you roll a fair six-sided die. You read off the number showing on the die, then flip that many fair coins.

(a) If the result of your die roll is i , what is the expected number of heads you see?

(b) What is the expected number of heads you see?

4 Number of Ones

Note 20

In this problem, we will revisit dice-rolling, except with conditional expectation. (*Hint*: for both of these subparts, the law of total expectation may be helpful.)

(a) If we roll a die until we see a 6, how many ones should we expect to see?

(b) If we roll a die until we see a number greater than 3, how many ones should we expect to see?