

Due: Saturday, 10/26, 4:00 PM
Grace period until Saturday, 10/26, 6:00 PM

Sundry

Before you start writing your final homework submission, state briefly how you worked on it. Who else did you work with? List names and email addresses. (In case of homework party, you can just describe the group.)

1 Count It!

Note 11

For each of the following collections, determine and briefly explain whether it is finite, countably infinite (like the natural numbers), or uncountably infinite (like the reals):

- (a) The integers which divide 8.
- (b) The integers which 8 divides.
- (c) The functions from \mathbb{N} to \mathbb{N} .
- (d) The set of strings over the English alphabet. (Note that the strings may be arbitrarily long, but each string has finite length. Also the strings need not be real English words.)
- (e) The set of finite-length strings drawn from a countably infinite alphabet, \mathcal{C} .
- (f) The set of infinite-length strings over the English alphabet.

2 Unprogrammable Programs

Note 12

Prove whether the programs described below can exist or not.

- (a) A program $P(F, x, y)$ that returns true if the program F outputs y when given x as input (i.e. $F(x) = y$) and false otherwise.
- (b) A program P that takes two programs F and G as arguments, and returns true if F and G halt on the same set of inputs (or false otherwise).

Hint: Use P to solve the halting problem, and consider defining two subroutines to pass in to P , where one of the subroutines always loops.

3 Kolmogorov Complexity

Note 12

Compressing a bit string x of length n can be interpreted as the task of creating a program of fewer than n bits that returns x . The Kolmogorov complexity of a string $K(x)$ is the length of an optimally-compressed copy of x ; that is, $K(x)$ is the length of shortest program that returns x .

- (a) Explain why the notion of the "smallest positive integer that cannot be defined in under 280 characters" is paradoxical.
- (b) Prove that for any length n , there is at least one string of bits that cannot be compressed to less than n bits, assuming that no two strings can be compressed to the same value.
- (c) Say you have a program K that outputs the Kolmogorov complexity of any input string. Under the assumption that you can use such a program K as a subroutine, design another program P that takes an integer n as input, and outputs the length- n binary string with the highest Kolmogorov complexity. If there is more than one string with the highest complexity, output the one that comes first lexicographically.
- (d) Let's say you compile the program P you just wrote and get an m bit executable, for some $m \in \mathbb{N}$ (i.e. the program P can be represented in m bits). Prove that the program P (and consequently the program K) cannot exist.

(Hint: Consider what happens when P is given a very large input n that is much greater than m .)

4 Probability Warm-Up

Note 13

- (a) Suppose that we have a bucket of 30 green balls and 70 orange balls. If we pick 15 balls uniformly out of the bucket, what is the probability of getting exactly k green balls (assuming $0 \leq k \leq 15$) if the sampling is done **with** replacement, i.e. after we take a ball out the bucket we return the ball back to the bucket for the next round?
- (b) Same as part (a), but the sampling is **without** replacement, i.e. after we take a ball out the bucket we **do not** return the ball back to the bucket.
- (c) If we roll a regular, 6-sided die 5 times. What is the probability that at least one value is observed more than once?

5 Five Up

Note 13

Say you toss a coin five times, and record the outcomes. For the three questions below, you can assume that order matters in the outcome, and that the probability of heads is some p in $0 < p < 1$, but *not* that the coin is fair ($p = 0.5$).

- (a) What is the size of the sample space, $|\Omega|$?

- (b) How many elements of Ω have exactly three heads?
- (c) How many elements of Ω have three or more heads?

For the next three questions, you can assume that the coin is fair (i.e. heads comes up with $p = 0.5$, and tails otherwise).

- (d) What is the probability that you will observe the sequence HHHTT? What about HHHHT?
- (e) What is the probability of observing at least one head?
- (f) What is the probability you will observe more heads than tails?

For the final three questions, you can instead assume the coin is biased so that it comes up heads with probability $p = \frac{2}{3}$.

- (g) What is the probability of observing the outcome HHHTT? What about HHHHT?
- (h) What about the probability of at least one head?
- (i) What is the probability you will observe more heads than tails?